

# Thinking Statistically about Analyzing Global Environmental Datasets

By Noel Cressie

Director, Program in Spatial Statistics and Environmental Sciences  
The Ohio State University  
(<http://www.stat.osu.edu/~sses>)

# Outline

Thinking Statistically: Regression

Thinking  
Statistically:  
Regression

MISR and its AOD Products

MISR and its AOD  
Products

Kriging

Kriging

Fixed Rank Kriging (FRK)

Fixed Rank Kriging  
(FRK)

FRK on a MISR Level 3 AOD Product

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking Statistically: Stratification and Aggregation

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

Conclusions

# Thinking Statistically

- ▶ Data are  $\{X_1, \dots, X_n\}$  at locations  $\mathbf{s}_1, \dots, \mathbf{s}_n$  on a given day: Summarize with a scatter diagram and  $(\bar{X}, S_X^2)$   
Example:  $X = \text{aerosol}$
- ▶ Data are  $\{Y_1, \dots, Y_n\}$  at *same* locations  $\mathbf{s}_1, \dots, \mathbf{s}_n$  on the *same* day: Summarize with a scatter diagram and  $(\bar{Y}, S_Y^2)$   
Example:  $Y = \text{precipitation}$

# Summaries

Global  
Environmental  
Datasets

N. Cressie

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products


Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions



$$\bar{X} = 2.6337 \quad S_X^2 = 0.2623$$



# Summaries

Global  
Environmental  
Datasets

N. Cressie

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products


Kriging

Fixed Rank Kriging  
(FRK)


FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions



$$\bar{X} = 2.6337 \quad S_X^2 = 0.2623$$



$$\bar{Y} = 3.6483 \quad S_Y^2 = 0.2976$$

Is  $Y$  related to  $X$ ?


Data are  $\{(X_1, Y_1), \dots, (X_n, Y_n)\}$  at locations  $\mathbf{s}_1, \dots, \mathbf{s}_n$  on a given day: Summarize with a scatter plot and a statistical regression relationship,

$$Y = \beta_0 + \beta_1 X + \text{error}$$

# Scatterplot of $Y$ versus $X$

Global  
Environmental  
Datasets

N. Cressie



$$Y = \beta_0 + \beta_1 X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product


Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Regression of $Y$ on $X$

Global  
Environmental  
Datasets

N. Cressie



$$Y = \beta_0 + \beta_1 X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Regression and Statistical Inference

- ▶ The fitted line  $Y = \hat{\beta}_0 + \hat{\beta}_1 X$  “fills in” missing values of  $Y$  for *any* given value of  $X$ .
- ▶ Try to imagine science without regression! We would have to let  $X$  vary on a very fine scale and measure each  $Y$  associated with these many  $X$ 's. Given  $X = X_0$ , we *infer*  $\hat{Y}_0 = \hat{\beta}_0 + \hat{\beta}_1 X_0$ .
- ▶ Are we certain that  $\hat{Y}_0 = Y_0$ , the true value? No, but we are 95% certain that

$$Y_0 \in (\hat{Y}_0 - 1.96\sigma_0, \hat{Y}_0 + 1.96\sigma_0),$$

where  $\sigma_0^2 = E(\hat{Y}_0 - Y_0)^2 =$  (well known formula)

# Regression and Statistical Inference

- ▶ The fitted line  $Y = \hat{\beta}_0 + \hat{\beta}_1 X$  “fills in” missing values of  $Y$  for *any* given value of  $X$ .
- ▶ Try to imagine science without regression! We would have to let  $X$  vary on a very fine scale and measure each  $Y$  associated with these many  $X$ 's. Given  $X = X_0$ , we *infer*  $\hat{Y}_0 = \hat{\beta}_0 + \hat{\beta}_1 X_0$ .
- ▶ Are we certain that  $\hat{Y}_0 = Y_0$ , the true value? No, but we are 95% certain that

$$Y_0 \in (\hat{Y}_0 - 1.96\sigma_0, \hat{Y}_0 + 1.96\sigma_0),$$

where  $\sigma_0^2 = E(\hat{Y}_0 - Y_0)^2 =$  (well known formula)

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Regression and Statistical Inference

Global  
Environmental  
Datasets

N. Cressie

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

- ▶ The fitted line  $Y = \hat{\beta}_0 + \hat{\beta}_1 X$  “fills in” missing values of  $Y$  for *any* given value of  $X$ .
- ▶ Try to imagine science without regression! We would have to let  $X$  vary on a very fine scale and measure each  $Y$  associated with these many  $X$ 's. Given  $X = X_0$ , we *infer*  $\hat{Y}_0 = \hat{\beta}_0 + \hat{\beta}_1 X_0$ .
- ▶ Are we certain that  $\hat{Y}_0 = Y_0$ , the true value? No, but we are 95% certain that

$$Y_0 \in (\hat{Y}_0 - 1.96\sigma_0, \hat{Y}_0 + 1.96\sigma_0),$$

where  $\sigma_0^2 = E(\hat{Y}_0 - Y_0)^2 =$  (well known formula)

# Spatial Regression

- ▶  $Y_0$  = aerosol at location  $\mathbf{s}_0$
- ▶  $\{X_j\}$  = aerosol values at *nearby* locations  $\{\mathbf{s}_j\}$
- ▶ Multiple regression:

$$Y_0 = \lambda_0 + \lambda_1 X_1 + \cdots + \lambda_m X_m + \text{error}$$

- ▶ Fill in missing aerosol values  $Y_0$  from observations  $\{X_j\}$ :

$$\hat{Y}_0 = \hat{\lambda}_0 + \hat{\lambda}_1 X_1 + \cdots + \hat{\lambda}_m X_m$$

and

$$\sigma_0^2 = E(\hat{Y}_0 - Y_0)^2$$

Recall

$$Y_0 \in (\hat{Y}_0 - 1.96\sigma_0, \hat{Y}_0 + 1.96\sigma_0)$$

with 95% probability

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation


Conclusions



# MISR Level 2 Aerosol Optical Depth (AOD) Coverage on April 1, 2002

Global  
Environmental  
Datasets

N. Cressie



Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

“Non-retrieval” occurs when radiance data are missing,  
clouds are present, or the algorithm fails

# MISR Level AOD (band 3, 558 nm)

## $0.5 \times 0.5$ degree on the globe

Global  
Environmental  
Datasets

N. Cressie

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products


Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions



$720 \times 360$  pixels in a half-degree by half-degree global map; Level 2 data are averaged within each pixel

# Spatial Statistical Model

- Observed variable

$$Z(\mathbf{s}) = Y(\mathbf{s}) + \varepsilon(\mathbf{s}); \quad \mathbf{s} \in D,$$

where  $\varepsilon(\cdot)$  is uncorrelated measurement error

- Hidden process with  $p$  regressors  $t_1(\mathbf{s}), \dots, t_p(\mathbf{s})$

$$Y(\mathbf{s}) = \mathbf{T}(\mathbf{s})' \alpha + \nu(\mathbf{s}); \quad \mathbf{s} \in D,$$

where  $\mathbf{T}(\mathbf{s}) \equiv (t_1(\mathbf{s}), \dots, t_p(\mathbf{s}))'$ , regression parameters  $\alpha$  are fixed but unknown, and  $\nu(\cdot)$  has mean zero

- Variance-covariance structure

$$\text{var}(\varepsilon(\mathbf{s})) = \sigma^2 \nu(\mathbf{s}), \quad \text{cov}(\nu(\mathbf{s}), \nu(\mathbf{t})) = \mathbf{C}(\mathbf{s}, \mathbf{t})$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Covariance Matrix of Data

Denote

$$\mathbf{V} \equiv \text{diag}(v(\mathbf{s}_1), \dots, v(\mathbf{s}_n))$$

$$\mathbf{C} \equiv \{C(\mathbf{s}_i, \mathbf{s}_j)\}$$

Define

$$\mathbf{\Sigma} \equiv \text{var}((Z(\mathbf{s}_1), \dots, Z(\mathbf{s}_n))')$$

Then

$$\mathbf{\Sigma} = \mathbf{C} + \sigma^2 \mathbf{V}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Kriging Equations

- The kriging predictor at location  $\mathbf{s}_0$  is

$$\begin{aligned}\hat{Y}(\mathbf{s}_0) &= \mathbf{T}(\mathbf{s}_0)' \hat{\alpha} + \mathbf{k}(\mathbf{s}_0)' (\mathbf{Z} - \mathbf{T} \hat{\alpha}) \\ &= \hat{\lambda}_1 Z(\mathbf{s}_1) + \cdots + \hat{\lambda}_n Z(\mathbf{s}_n)\end{aligned}$$

with

$$\mathbf{T} \equiv (\mathbf{T}(\mathbf{s}_1), \dots, \mathbf{T}(\mathbf{s}_n))', \quad \hat{\alpha} = (\mathbf{T}' \boldsymbol{\Sigma}^{-1} \mathbf{T})^{-1} \mathbf{T}' \boldsymbol{\Sigma}^{-1} \mathbf{Z},$$

$$\text{and } \mathbf{k}(\mathbf{s}_0)' = \mathbf{c}(\mathbf{s}_0)' \boldsymbol{\Sigma}^{-1}$$

- Now let  $\mathbf{s}_0$  vary to create a map of  $\hat{Y}(\cdot)$

# Kriging Equations, ctd.

- ▶ The root-mean-squared prediction error (kriging standard error) of  $\hat{Y}(\mathbf{s}_0)$  is:

$$\sigma_k(\mathbf{s}_0) = \left\{ \mathbf{C}(\mathbf{s}_0, \mathbf{s}_0) - \mathbf{k}(\mathbf{s}_0)' \mathbf{\Sigma} \mathbf{k}(\mathbf{s}_0) \right. \\ \left. + (\mathbf{T}(\mathbf{s}_0) - \mathbf{T}' \mathbf{k}(\mathbf{s}_0))' (\mathbf{T}' \mathbf{\Sigma}^{-1} \mathbf{T})^{-1} \right. \\ \left. (\mathbf{T}(\mathbf{s}_0) - \mathbf{T}' \mathbf{k}(\mathbf{s}_0)) \right\}^{1/2}$$

- ▶  $Y(\mathbf{s}_0) \in (\hat{Y}(\mathbf{s}_0) - 1.96\sigma_k(\mathbf{s}_0), \hat{Y}(\mathbf{s}_0) + 1.96\sigma_k(\mathbf{s}_0))$  with 95% probability.

# Kriging of Very Large Data Sets

- ▶ The bottleneck for kriging computations is inverting the  $n \times n$  covariance matrix  $\Sigma$
- ▶ Directly inverting  $\Sigma$  becomes very slow, even infeasible, when sample size  $n$  is large (e.g., when  $n > 3,000$ )
- ▶ For the MISR Level 3 global AOD data on half-degree resolution,  $n$  could be as much as  $720 \times 360 = 295,200$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Fixed Rank Kriging (FRK)

- ▶  $\Sigma = \mathbf{C} + \sigma^2 \mathbf{V}$
- ▶ Assume  $\mathbf{C}_{n \times n}$  is of the form

$$\mathbf{C} = \mathbf{S}_{n \times r} \mathbf{K}_{r \times r} \mathbf{S}_{r \times n}' ,$$

with  $r$  fixed basis functions,

$\{\mathbf{S}_j(\cdot) : j = 1, \dots, r\} \equiv \mathbf{S}(\cdot)$ , and  $r \ll n$

- ▶ We have derived explicit forms of  $\Sigma^{-1}$ , and hence of the kriging solutions, when  $n$  is very large. Inversion of  $r \times r$  matrices and  $n \times n$  diagonal matrices are involved
- ▶ References: Cressie and Johannesson (2006, 2008) and Shi and Cressie (2007)



# Fixed Rank Kriging Equations

$$\Sigma^{-1} = (\sigma^2 \mathbf{V})^{-1} - (\sigma^2 \mathbf{V})^{-1} \mathbf{S} (\mathbf{K}^{-1} + \mathbf{S}' (\sigma^2 \mathbf{V})^{-1} \mathbf{S})^{-1} \mathbf{S}' (\sigma^2 \mathbf{V})^{-1}$$

$$\begin{aligned}\hat{Y}(\mathbf{s}_0) &= \mathbf{T}(\mathbf{s}_0)' \hat{\alpha} + \mathbf{S}(\mathbf{s}_0)' \mathbf{K} \mathbf{S}' \Sigma^{-1} (\mathbf{Z} - \mathbf{T} \hat{\alpha}) \\ \hat{\alpha} &= (\mathbf{T}' \Sigma^{-1} \mathbf{T})^{-1} \mathbf{T}' \Sigma^{-1} \mathbf{Z}\end{aligned}$$

$$\begin{aligned}\sigma_k(\mathbf{s}_0) &= \{ \mathbf{S}(\mathbf{s}_0)' \mathbf{K} \mathbf{S}(\mathbf{s}_0) - \mathbf{S}(\mathbf{s}_0)' \mathbf{K} \mathbf{S}' \mathbf{K} \mathbf{S}(\mathbf{s}_0) \\ &\quad + (\mathbf{T}(\mathbf{s}_0) - \mathbf{T}' \Sigma^{-1} \mathbf{S} \mathbf{K} \mathbf{S}(\mathbf{s}_0))' (\mathbf{T}' \Sigma^{-1} \mathbf{T})^{-1} \\ &\quad (\mathbf{T}(\mathbf{s}_0) - \mathbf{T}' \Sigma^{-1} \mathbf{S} \mathbf{K} \mathbf{S}(\mathbf{s}_0)) \}^{1/2}\end{aligned}$$

# Estimating $\mathbf{K}$

- ▶ Given the empirical, binned covariance matrix  $\hat{\Sigma}_M$ , we find  $\hat{\mathbf{K}}$  and  $\hat{\sigma}^2$  that minimize the Frobenius norm between  $(\mathbf{S}_M \mathbf{K} \mathbf{S}_M' + \sigma^2 \mathbf{V}_M)$  and  $\hat{\Sigma}_M$
- ▶ Frobenius norm between two matrices:

$$\|\mathbf{A} - \mathbf{B}\|^2 = \text{tr}((\mathbf{A} - \mathbf{B})'(\mathbf{A} - \mathbf{B}))$$

(We use a weighted version of the Frobenius norm to estimate  $\mathbf{K}$  and  $\sigma^2$ )

# Estimating $\mathbf{K}$ , ctd.

- By minimizing the Frobenius norm, we obtain

$$\hat{\mathbf{K}} = \mathbf{R}^{-1} \mathbf{Q}' (\hat{\boldsymbol{\Sigma}}_M - \hat{\sigma}^2 \mathbf{V}_M) \mathbf{Q} (\mathbf{R}^{-1})',$$

where  $\mathbf{Q}$  and  $\mathbf{R}$  are obtained from the  $Q$ - $R$  decomposition of  $\mathbf{S}_M$


- Now we have all we need for FRK!

# MISR Level 3 AOD Data, APR 01-16, 2001


Global  
Environmental  
Datasets

N. Cressie

Histogram of AOD



Histogram of  $\log(\text{AOD})$



Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

$$Z(\mathbf{s}) = \log(\text{AOD}(\mathbf{s}))$$




Region of Interest ( $D$ ):  $\text{lat}(-20, 45)$ ,  $\text{lon}(-140, 12)$

# Region of Interest

Total number of pixels: 32,768 ( $128 \times 256$ )

Pixels observed:  $n = 25,897$  (79% of total no. pixels)

Observed log(AOD), APR 01-16, 2001



Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product


Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions



# Residuals from OLS

$$Z_R(\mathbf{s}_0) \equiv Z(\mathbf{s}_0) - \mathbf{t}(\mathbf{s}_0)' \hat{\alpha}_{OLS}$$



These residuals are used to fit  $\mathbf{K}$  in FRK

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions




# FRK Results


Global  
Environmental  
Datasets

N. Cressie


Observed log(AOD)



FRK Prediction of log(AOD)



FRK SE of log(AOD)



Components of  $\mathbf{S}(\cdot)$  are  $W$ -wavelets at finer resolutions

Fixed rank of  $\mathbf{K} = r =$  no. of basis functions = 300

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Computation Time

Computation is done in Matlab on a Pentium 4 dual core 3.0Ghz, 2GB memory Linux Machine. Time unit: seconds

$p$	$r$	Fit $\mathbf{K}$ , $\sigma^2$	Prediction
32	300	1895.3	156.0
32	400	2536.8	278.7
64	300	1894.4	170.2
64	400	2536.6	287.6

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

- ▶ FRK: Computations are  $O(n)$  as compared to the usual kriging algorithms, where they are  $O(n^3)$ . *All*  $n$  data are used to produce the **optimal** linear predictor
- ▶ Multivariate: The same ideas could be used to combine MISR and MODIS data to obtain an optimal predictor of AOD (**data fusion**)
- ▶ **Space-time**: Occasional, off-line estimation of **K** would be possible for fast, optimal updating of the map at regular time intervals. As well as current AOD values, past values could be used to improve on FRK (i.e., smaller MSPEs); my IWGGMS-5 paper at Caltech on 6/25/08 addresses this problem

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Epilogue: Still Thinking Statistically

Global  
Environmental  
Datasets

N. Cressie

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

- ▶ Regression depends as much on what  $X$  is chosen to regress on  $Y$  as what other variable was *not* chosen
- ▶ The variable  $A$  = cloud microphysical properties (say), potentially influences the (regression) relationship between  $Y$  = precipitation, and  $X$  = aerosol

# Stratify (on variable $A$ )

$$\begin{aligned}\text{Stratum 1} &: \{\mathbf{s}_i: i \in A_1\} \\ \text{Stratum 2} &: \{\mathbf{s}_i: i \in A_2\} \\ \text{Stratum 3} &: \{\mathbf{s}_i: i \in A_3\} \\ A_1 \cup A_2 \cup A_3 &= \{1, \dots, n\}\end{aligned}$$

Summarize with a scatter plot that highlights the three different strata, and three statistical regression relationships. In region  $k$  ( $k = 1, 2, 3$ ):

$$Y = \beta_{0,k}^{(A)} + \beta_{1,k}^{(A)}X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Scatterplot of $Y$ versus $X$



$$Y = \beta_0 + \beta_1 X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions


# Stratification on Variable A



For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(A)} + \beta_{1,k}^{(A)} X + \text{error}$$

# Stratification on Variable A



$$\hat{\beta}_{1,1}^{(A)} > 0$$

$$\hat{\beta}_{1,2}^{(A)} > 0$$

$$\hat{\beta}_{1,3}^{(A)} > 0$$

For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(A)} + \beta_{1,k}^{(A)}X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)


FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions



# Scatterplot of $Y$ versus $X$



$$Y = \beta_0 + \beta_1 X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Stratification on Variable $B$



For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(B)} + \beta_{1,k}^{(B)} X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Stratification on Variable $B$



$$\hat{\beta}_{1,1}^{(B)} < 0$$

$$\hat{\beta}_{1,2}^{(B)} < 0$$

$$\hat{\beta}_{1,3}^{(B)} > 0$$

For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(B)} + \beta_{1,k}^{(B)}X + \text{error}$$

# Aggregate (on variable A)

In Stratum  $k$  ( $k = 1, \dots, K$ ): Summarize with aggregated data  $(\bar{X}_k, \bar{Y}_k)$

This is done all the time!

Derived (aggregated) data:  $\{(\bar{X}_1, \bar{Y}_1), \dots, (\bar{X}_K, \bar{Y}_K)\}$  for Stratum 1, ..., Stratum  $K$  on a given day.

Summarize with a scatter plot and a statistical regression relationship,

$$\bar{Y} = \alpha_0 + \alpha_1 \bar{X} + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions


# Stratification on Variable A



For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(A)} + \beta_{1,k}^{(A)} X + \text{error}$$

# Stratification on Variable A



$$\hat{\beta}_{1,1}^{(A)} > 0$$

$$\hat{\beta}_{1,2}^{(A)} > 0$$

$$\hat{\beta}_{1,3}^{(A)} > 0$$

For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(A)} + \beta_{1,k}^{(A)}X + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging


Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions


# Stratification on Variable A



For 3 strata ( $k = 1, 2, 3$ )

$$Y = \beta_{0,k}^{(A)} + \beta_{1,k}^{(A)} X + \text{error}$$

# Aggregation Using Variable A



$$\overline{Y} = \alpha_0^{(A)} + \alpha_1^{(A)} \overline{X} + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)


FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions



# Aggregation Using Variable A



$$\hat{\alpha}_1^{(A)} < 0$$

$$\bar{Y} = \alpha_0^{(A)} + \alpha_1^{(A)}\bar{X} + \text{error}$$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Conclusions

- ▶ Regression **fills gaps** in noisy data with known mean squared prediction error
- ▶ **Kriging** (spatial regression) fills gaps in noisy spatial data with known kriging variance
- ▶ **Very large spatial datasets** can be kriged with a flexible family of spatial covariance functions
- ▶ Regression requires care when stratifying and aggregating. Solution: Build **conditional probability models**:  $[Y|X, A]$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Conclusions

- ▶ Regression **fills gaps** in noisy data with known mean squared prediction error
- ▶ **Kriging** (spatial regression) fills gaps in noisy spatial data with known kriging variance
- ▶ Very large spatial datasets can be kriged with a flexible family of spatial covariance functions
- ▶ Regression requires care when stratifying and aggregating. Solution: Build **conditional probability models**:  $[Y|X, A]$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Conclusions

- ▶ Regression **fills gaps** in noisy data with known mean squared prediction error
- ▶ **Kriging** (spatial regression) fills gaps in noisy spatial data with known kriging variance
- ▶ **Very large spatial datasets** can be kriged with a flexible family of spatial covariance functions
- ▶ Regression requires care when stratifying and aggregating. Solution: Build **conditional probability models**:  $[Y|X, A]$

Thinking  
Statistically:  
Regression

MISR and its AOD  
Products

Kriging

Fixed Rank Kriging  
(FRK)

FRK on a MISR  
Level 3 AOD  
Product

Still Thinking  
Statistically:  
Stratification and  
Aggregation

Conclusions

# Conclusions

- ▶ Regression **fills gaps** in noisy data with known mean squared prediction error
- ▶ **Kriging** (spatial regression) fills gaps in noisy spatial data with known kriging variance
- ▶ **Very large spatial datasets** can be kriged with a flexible family of spatial covariance functions
- ▶ Regression requires care when stratifying and aggregating. Solution: Build **conditional probability models**:  $[Y|X, A]$